




Apr 4th, 4:45 PM - 5:00 PM

**The effects of diluted bitumen (dilbit) exposure during embryonic development on the future swimming performance and metabolic and ionic recovery post-exercise in sockeye salmon (*Oncorhynchus nerka*)**

Feng Lin  
*Simon Fraser Univ., Canada, fla44@sfu.ca*

Chris Kennedy  
*Simon Fraser Univ., Canada, ckennedy@sfu.ca*

Follow this and additional works at: <https://cedar.wvu.edu/ssec>

 Part of the [Fresh Water Studies Commons](#), [Marine Biology Commons](#), [Natural Resources and Conservation Commons](#), and the [Terrestrial and Aquatic Ecology Commons](#)

---

Lin, Feng and Kennedy, Chris, "The effects of diluted bitumen (dilbit) exposure during embryonic development on the future swimming performance and metabolic and ionic recovery post-exercise in sockeye salmon (*Oncorhynchus nerka*)" (2018). *Salish Sea Ecosystem Conference*. 116.  
<https://cedar.wvu.edu/ssec/2018ssec/allsessions/116>

This Event is brought to you for free and open access by the Conferences and Events at Western CEDAR. It has been accepted for inclusion in Salish Sea Ecosystem Conference by an authorized administrator of Western CEDAR. For more information, please contact [westerncedar@wvu.edu](mailto:westerncedar@wvu.edu).

# The effects of diluted bitumen on the embryonic development and swimming performance of sockeye salmon (*Oncorhynchus nerka*)




Feng Lin and Chris Kennedy  
Simon Fraser University, British Columbia, Canada  
30<sup>th</sup> Salish Sea Ecosystem Conference, April 4-6, 2018






# Background



# Knowledge gap about dilbit toxicity

- Dilbit, a non-conventional crude oil
  - ↓ BTEX
  - ↓ Total polycyclic aromatic hydrocarbons (PAHs)
  - ↑ 3~5 ringed PAHs in proportions
    - Most toxic and bioavailable to aquatic life
- Different environmental behaviour when spilled
  - High density and viscosity
  - High potential of sinking



Processes affecting the composition, amount, and behaviour of diluted bitumen



# Knowledge gap about dilbit toxicity

## ➤ Dilbit toxicity to fish species remains unclear

- Very few studies on dilbit toxicity
- Not enough empirical data

## ➤ Studies using other crude oil blend or its components (e.g. BTEX, PAHs)

- Generalization of risk?



Sockeye salmon



Japanese medaka




Fathead minnow




Inland silverside

### **BTEX**





### **PAHs**



# Study objectives

- Acute and sublethal toxicity of dilbit on early life stage (ELS) Pacific salmonids
  - Mortality
  - % Deformity
  - Swimming ability
  - Biochemistry
- Latent effect in formally exposed survivors?
- Environmentally relevant exposure method





# Exposure apparatus



Siporax® ceramic beads



Dilbit


Pre-soaked for 48h



- Generate water soluble fractions (WSFs) of dilbit
- Initial spike of total PAHs at 3.5, 16.4, 66.7  $\mu\text{g/L}$
- Beads re-soaked every 21 days

# Experiment part 1

- Fertilized sockeye salmon embryos exposed to 3 concentrations of WSFs
  - 3.5, 16.4, 66.7  $\mu\text{g/L}$  total PAHs
  - Control at 0.002  $\mu\text{g/L}$
- At 50% yolk sac absorption
  - Deformity assessment
- At swim-up stage
  - Burst swimming ability
  - Body biochemistry




Embryo and alevin stage




Swim-up stage fry  
(no external yolk sac)

# Results



Values are means  $\pm$  s.e.m (n = 960). One-factor ANOVA and Tukey HSD test ( $p < 0.05$ ).



The stacked cumulative mortality (%) of fish across treatments (n = 960) at three different developmental stages




	Control	3.5 µg/L	16.4 µg/L	66.7 µg/L
Deformity (%)				
Yolk sac edema	2.0±1.0 <sup>c</sup>	6.0±1.0 <sup>b</sup>	15.0±4.0 <sup>a</sup>	12.0±2.0 <sup>a</sup>
Pericardiac edema	1.0±0	0	4.0±2.0	1.0±0
Craniofacial	1.0±0	3.0±2.0	3.0±1.0	5.0±2.0
Skeletal	0	1.0±0	1.0±0	1.0±0
Finfold	0	1.0±0	0	0
Fish with at least 1 type of deformity	2.0±1.0 <sup>c</sup>	8.0±1.0 <sup>b</sup>	20.0±2.0 <sup>a</sup>	17.5 ±0.5 <sup>a</sup>


Values are means ± s.e.m (n = 200). One-factor ANOVA and Tukey HSD test (p<0.05)




Skeletal deformity



Craniofacial deformity



Presence of edema





Finfold deformity



# Results




Swim-up stage fry



Values are means  $\pm$  95% confidence interval (n = 20). One-factor ANOVA and Tukey HSD test (p<0.05).

# Results



Swim-up stage fry


- Unchanged carbohydrate storage
- Increased lipid storage
- Decreased soluble protein

Values are means  $\pm$  95% confidence interval (n = 20). One-factor ANOVA and Tukey HSD test (p<0.05).




Swim-up fry

- $U_{burst}$  swimming speed --- measure of burst ability
  - Involved in food capture and predation avoidance



Loligo swimming tunnel used for  $U_{burst}$  test



Values are means  $\pm$  95% confidence interval (n = 12). One-factor ANOVA and Tukey HSD test ( $p < 0.05$ ).



Exposed swim-up fry





Yearling parr



Pre-smolt stage

Reared in clean water for 8 months.  $U_{burst}$  swimming speed tested at 1, 3, 6, or 8 months in clean water








Exposed swim-up fry



Yearling parr






Pre-smolt stage



# Pre- and post- exercise biochemistry



Values are means  $\pm$  95% confidence interval (n = 12).  
One-factor ANOVA and Tukey HSD test (p<0.05).



- Indicators of stress level
- Spikes in circulating cortisol, [glucose], [lactate] after exhaustive exercise



# Pre- and post- exercise biochemistry



➤ Indicators of water-ion homeostasis


➤ Reduction in circulating plasma [Na<sup>+</sup>] and [Cl<sup>-</sup>] after exhaustive exercise

Values are means  $\pm$  95% confidence interval (n = 12). One-factor ANOVA and Tukey HSD test (p<0.05).



# Conclusions

## Exposure to WSFs of dilbit:



- Latent effect
- Signs of recovery after 8M?

- $\uparrow$  Mortality
- $\uparrow$  % Deformity

- $\downarrow U_{burst}$  swimming speed
- Altered biochemistry



## Experiment part 2 (ongoing)

- Exposed juvenile sockeye salmon exposed to 3 concentrations of WSFs
  - 3 exposure length: 24h, 96h, or 14d
- $U_{burst}$  and body biochemistry
  - Exercise recovery ability
  - Stress response



Pre-smolt stage



Exposure set-up

# Thank you for listening!

## Acknowledgements:



Fisheries and Oceans  
Canada


Pêches et Océans  
Canada

Funding support by the National  
Contaminant Advisory Group

**SFU**

**SIMON FRASER UNIVERSITY**  
ENGAGING THE WORLD

Supervision by Dr. Chris Kennedy  
Animal care staff at SFU for extensive  
fish care





Water accommodated fractions (WAFs)